# **COSMETIC SURGERY OF LAS VEGAS**

Patient:



#### COSMETIC SURGERY OF LAS VEGAS

DR SAMIR PANCHOLI

### **Facial Enhancement - Post Surgery Care**

Face lift, Neck Lift, Mid-face Lift, Forehead and Brow Lift

**Shopping:** Obtain these items before surgery.

- Prescriptions
  - Supplies:
    - Ice pack or Frozen peas and Ziploc bags.
    - Q-Tips
    - Hydrogen Peroxide
    - Antibiotic Ointment
    - A New Soft Bristle Toothbrush
- Food Day 1: Simple, easy to digest foods (toast, crackers, Ensure or Boost, pudding, soup, etc.) Week 1 and 2: Soft diet.
  - Week 3 and forward: Slowly progress to a normal diet.

### Dressings/Bandages:

- Your head wrap will be removed the next day during your appointment.
- Starting the day after surgery
  - Face or Neck Lift Patients: Wear your neck support strap after surgery as follows:
    - First week 24 hours a day
    - Second week At least 12 hours a day
    - Third and Fourth weeks At night
    - You may remove your neck support strap when you take a shower.
- You may shower daily starting 24 hours after surgery.
  - Do not bathe.
  - The surgical tape / foam may get wet, but avoid soaking the tape / foam in water.
  - Towel dry the surgical tape / foam first and then use a hairdryer on a cool air setting to complete the drying process. Do not use the warm or heated hairdryer settings

### Sutures:

- Your sutures are dissolvable
- Any staples in hair bearing areas (if applicable) will be removed at your one-week appointment.

### Wound Care:

- Starting the first day *after* surgery, clean your incisions as follows:
  - Keep your surgical tape / foam in place.
  - Wash your hair gently everyday.
  - After your shower
    - Gently dab all incisions using a Q-tip soaked in diluted hydrogen peroxide (mixed with 50% water).
    - Then apply a thin layer of antibiotic ointment to <u>all</u> incisions.
    - Dry the surgical tape / foam as described above using a hairdryer on a cool air setting.
    - Mid Face Lift patients with oral incisions Perform the following three times a day:
    - Carefully brush your teeth
      - Rinse with warm salty water.

## **COSMETIC SURGERY OF LAS VEGAS**

#### Patient:

#### Activity:

- Up to 6 weeks:
  - Limit neck turning
    - Avoid rapidly turning your head
- Up to 3 weeks:
  - Do nothing that raises your heart rate or blood pressure including but not limited to:
  - Lifting
  - Exercise
- Up to 2 weeks:
  - Avoid heavy exercise or lifting
- Up to 1 week:
  - Avoid driving while still taking prescription pain medication
  - Sleep on your back and avoid lying directly on the side of your face.
  - Sleep with your head elevated 30 to 45 degrees with several pillows to help decrease swelling.
  - Maintain a soft diet and avoid heavy chewing and prolonged talking
- <u>Up to 24 48 hours:</u>
  - Apply ice pack / cold pack / Frozen peas in Ziploc bags to your face (do not apply directly to skin)
  - Avoid all nicotine products (including second hand smoke) for at least 4 weeks.
  - Avoid drinking alcoholic beverages to minimize bleeding.
  - Do not lift anything over 10 pounds.

#### **Special Instructions:**

- Swelling, bruising, and some bloody drainage are normal for the first 2 weeks.
- Be aware of a possible hematoma by noticing:
  - Usually one side of your face begins to experience:
    - Extreme swelling
    - Extreme hardness
    - Extreme pain
  - Notify Dr. Pancholi immediately if the above occurs.
- If Drains were used:
  - They will be left in place until drainage is less that 30 cc over a 24-hour period.
  - Antibiotics must be taken as long as drains are in place. If you run out, call us for a refill.
  - Numbness, tingling, or sharp sensations is expected for the first few months.
- Do not drive any vehicles or operate and heavy machinery:
  - 36 hours following your surgery.
  - 4 hours following your last dose of pain medication.
  - 8 hours following your last dose of sleep medication.
- Prescriptions:

•

Patient

- If you have a sensitive stomach, take your antibiotic with yogurt.
- If your pain medicine is making you sick:
  - Lower the dose.
  - Take it with food.
  - Notify Dr. Pancholi.

	Date:
Relation	
to Patient	Date:
	Date:
	Date:
	Relation to Patient